



# topcote

**100% coated fertiliser**

- Excellent price/quality ratio
- Formula for nurseries
- Formula for horticulturists
- Versatility and safety

## Description of the product

FERTIL has a strong know-how in fertilisers; we can offer you a new coated fertiliser, with a very good price/quality ratio.

Two balances have been chosen :

- formula 11-11-27 ,release duration about 5-6 months,
- formula 17-11-11 , release duration about 8-9 months.

Granulometry : 2 – 2,5 mm.

## COMPONENTS

- formula 11-11-27 : Nitrogen (N) 11% with 8% of nitric N, 3% of amoniacal N, phosphorical Anhydride ( $P_2O_5$ ) 11% with 9,9% soluble in water ; Oxyde de Potassium ( $K_2O$ ) 27% soluble in water. Copper (Cu) 0.011%, manganese (Mn) 0.015%, molybdenum (Mo) 0.002%, zinc (Zn) 0.015%, Iron (Fe) 0.1%
- formula 17-11-11 : Nitrogen (N) 17% with 5% nitric N, 7% amoniacal N et 5% urea N; phosphorical Anhydride ( $P_2O_5$ ) 11% with 9.9% soluble in water. Oxyde of potassium ( $K_2O$ ) 11%. Bore (B) 0.02%, Copper (Cu) 0.04%, manganese (Mn) 0.05%, molybdenum (Mo) 0.007%, zinc (Zn) 0.05%, Iron (Fe) 0.3%

## Packaging

- 25 kg.bag
- 40 bags pallets (1 ton).



## Use

### RECOMMENDATIONS AND DOSIS FOR A NON-FERTILISED SUBSTRATE

Balance Type	Recomendations	Usage Doses	Ref.
11-11-27 6H	<ul style="list-style-type: none"><li>• Bed plants</li><li>• Flower pots</li><li>• Chrysanthemums</li><li>• Surfacing</li></ul>	<ul style="list-style-type: none"><li>• 2 à 4 kg/m<sup>3</sup></li><li>• 1 à 2 g/L</li></ul>	735.00
17-11-11 8P	<ul style="list-style-type: none"><li>• Nurseries plants in containers (open air)</li><li>• Surfacing</li></ul>	<ul style="list-style-type: none"><li>• 3 à 5 kg/m<sup>3</sup></li><li>• 1 à 3 g/L</li></ul>	739.00

### USE RECOMANDATIONS

When using TOPCOTE, you have to keep in mind the needs of th cultivated species and their method of cultivation. The general recommendations presented above must be adapted to each situation as well.

FERTIL can't be held responsible for damages linked to a bad use of the product.

The release starts since TOPCOTE is put in a humid environment Indeed, you shouldn't incorporate TOPCOTE more than one or two weeks in advance in your breeding ground.

[www.fertil.fr](http://www.fertil.fr)